

Self-Gravitating Strings and String/Black Hole Correspondence

Ramzi R. Khuri*

*Department of Natural Sciences, Baruch College, CUNY
17 Lexington Avenue, New York, NY 10010[†]*

*Graduate School and University Center, CUNY
365 5th Avenue, New York, NY 10036*

*Center for Advanced Mathematical Sciences
American University of Beirut, Beirut, Lebanon ***

Abstract

In a recent essay, we discussed the possibility of using polymer sizing to model the collapse of a single, long excited string to a black hole. In this letter, we apply this idea to bring further support to string/black hole correspondence. In particular, we reproduce Horowitz and Polchinski's results for self-gravitating fundamental strings and speculate on the nature of the quantum degrees of freedom of black holes in string theory.

October, 1999

* e-mail: khuri@gursey.baruch.cuny.edu. Supported by NSF Grant 9900773 and by PSC-CUNY Award 669663.

[†] Permanent address.

** Associate member.

Susskind's postulate of string-black hole correspondence [1] represents the most promising attempt to understand the physical basis of quantum gravity from string theory. In its most basic form, it suggests a one to one correspondence between fundamental string states and black hole states.

This correspondence was clarified further in [2], where it was argued that by adiabatically increasing the string coupling constant g a string state would turn into a black hole at $g = g_c \sim N^{-1/4}$, where $N \gg 1$ is the level number of a long, fundamental string state. The string coupling takes this critical value precisely when the Schwarzschild radius, R_S , becomes of the order of the string scale, l_s . For couplings below g_c , the picture of a string state prevails, while for coupling above g_c the black hole picture prevails. At this critical transition point, the string entropy makes a smooth transition into the Bekenstein-Hawking black hole area entropy law [3].

The authors of [2] took this correspondence a step further in [4], using a thermal scalar field theory formalism to study the size of the string state as it collapses from its initial random walk [5] form into a black hole. It was then argued in [6] that these results should also arise using the methods of polymer physics¹. More recently, the results of [2] were also reproduced in [8], and where some of the physical issues relating to the collapsing string were further clarified.

In this letter, we reproduce the results of [4] using polymer methods, making precise the scaling arguments of [6] and thereby giving further support to string-black hole correspondence. We also discuss the implications of these results to the understanding of the underlying degrees of freedom of quantum black holes in string theory.

Following [4], we consider a long, self-gravitating string at level N and adiabatically increase the coupling g until the string collapses into a black hole. As noted in [4], the string size at zero coupling (the free string) is initially given by $R_{RW} \sim N^{1/4}l_s$, where l_s is the string scale. The letters "RW" denote "Random Walk", as the free string represents a

¹ The concepts of the "string bit" and the "polymer string" were first discovered by Thorn [7].

random walk [5] with $n = N^{1/2}$ steps (or string “bits” [7]). The total length of the string is given by $L = nl_s$.

As argued in [6], this configuration may be represented as a random walk polymer chain with self-interaction. There are n steps, each of length l , with $\vec{r}_i$ representing the position of the chain after the i th step. Gravitational self-interactions start to become significant once $g \sim g_0 \sim n^{(d-6)/4}$. This can be seen immediately by noting that the attractive potential

$$U = \sum_{i \neq j} \frac{g^2}{|\vec{r}_i - \vec{r}_j|^{d-2}} \sim \frac{n^2 g^2}{n^{(d-2)/2} l_s^{d-2}} \sim g^2 n^{-(d-6)/2} (l_s^{2-d}) \quad (1)$$

becomes non-negligible in string units only when g approaches g_0 . Here we have used the fact that the mean square average of the distance between steps (the radius of gyration [9]) is proportional to the random walk size, $R_{RW} \sim n^{1/2} l_s$. We then replaced accordingly in the summation, up to a factor of the order unity (the precise numerical coefficient is not necessary for our purposes).

As in [6], we follow the calculation of [10] but instead of a self-avoiding delta-function interaction, we consider a self-attracting potential U as above. For simplicity, we restrict ourselves initially to the case of $D = 4$ spacetime dimensions (or $d = 3$ space dimensions).

The system is described by the generalized Hamiltonian

$$\beta H = \frac{3}{2l} \int_0^L ds \left(\frac{\partial \vec{R}(s)}{\partial s} \right)^2 + g^2 l \int_0^L \int_0^L ds ds' \frac{1}{|\vec{R}(s) - \vec{R}(s')|}, \quad (2)$$

where $\vec{R}(s)$ is the position vector of the chain at arc-length s ($0 \leq s \leq L$). Following [10], we use the Feynman variational procedure for the free energy of the chain. The trial Hamiltonian is given by

$$\beta H_0 = \frac{3}{2l_s} \int_0^L ds \left(\frac{\partial \vec{R}(s)}{\partial s} \right)^2 + Q, \quad (3)$$

where

$$Q = \frac{q^2}{6l_s} \int_0^L ds \vec{R}(s)^2, \quad (4)$$

where q is the variational parameter. The free energy of the chain is given by

$$\exp(-\beta F) = \int D\vec{R} \exp(-\beta H_0 + X + Q), \quad (5)$$

where

$$X = \frac{g^2}{l_s} \int_0^L \int_0^L ds ds' \frac{ds ds'}{|\vec{R}(s) - \vec{R}(s')|}, \quad (6)$$

and where q is a variational parameter. F is to be extremized with respect to q . Straightforward calculations similar to those of [10] yield the free energy

$$\beta F = \beta H_0 - Q - X = \frac{qL}{4} - \epsilon \frac{g^2 L^2}{l} \left(\frac{q}{l_s} \right)^{1/2}, \quad (7)$$

where ϵ is a number of the order unity. It is also straightforward to show that the size of the polymer, the average mean square end-to-end distance of the chain, is given by

$$R^2 = \frac{l_s}{q_0} (1 - \exp(-q_0 L)), \quad (8)$$

where q_0 is the value of q obtained by varying (7) with respect to q and is determined up to a factor of $O(1)$. The variation of (7) yields $q_0 L \sim g^4 n^3$, which implies

$$R^2 \simeq \frac{l_s^2}{g^4 n^3} (1 - \exp(-g^4 n^3)). \quad (9)$$

This finding makes precise the arguments of [6], and agrees with the results of [4]. In particular, note that for $g \ll g_0$, $R^2 \sim n l_s^2$, which is the random walk/free string result, while for $g_0 < g < g_c$, $R \sim l_s / (g^2 n)$, which agrees with the calculation of [4].

The above results can be generalized in a straightforward manner to the case of arbitrary spacetime dimension $D = d + 1$ (note, however, that the calculations are not reliable for $d > 4$). In this case, the free energy is given by

$$\beta F = \frac{qL}{4} - \epsilon \frac{g^2 L^2}{l_s^{4-d}} \left(\frac{q}{l} \right)^{\frac{1}{2}(d-2)}, \quad (10)$$

where again ϵ is a number of the order unity. The size of the polymer is again given by (8). The variation for general $d \neq 4$ yields $q_0 L \sim (g^4 n^{6-d})^{1/(4-d)}$, which again predicts a

transition from the random walk at $g_0 \sim n^{(d-6)/4}$, although the nature of the transition is somewhat more obscure [4,8].

The special case of $d = 4$ ($D = 5$) deserves some attention. Here $F \sim qL(1 - \epsilon g^2 n)$, so that for $g < g_c \sim n^{-1/2}$, F is minimized at $q = 0$, corresponding to the random walk, while for $g > g_c$, F is minimized for $q \rightarrow \infty$, so that $R^2 \rightarrow 0$, corresponding to a rapid collapse at the transitional coupling, which coincides in this case with the critical coupling: $g_0 = g_c$. This is again in agreement with the results of [4] and [8]. Note that the five-dimensional case also corresponds to the expected behaviour for a string state winding around the fifth dimension, and whose collapse results in a charged black hole.

Further agreement with the results of [4] can be seen by considering the temperature. To leading order, $T = T_H = 1/\beta = 1/l_s$, where T_H is the Hagedorn temperature, so we will compute the leading order correction. Again, for simplicity, let us focus on the case of $D = 4$ ($d = 3$). Then at $q = q_0$, $\beta F = \beta(E - TS) \sim g^4 n^3$. Using $E = M \sim n/l_s$, we obtain, to leading order, $TS \sim n/l_s(1 - \epsilon g^4 n^2)$. Since $S \sim n$ throughout the adiabatic collapse, this gives $T \sim (1 - \epsilon g^4 n^2)/l_s$. This again reproduces the results of [4], namely that $\beta - \beta_H \sim g^4 n^2 l_s \sim g^4 M^2 l_s^3 \sim \alpha'^2 g^4 M^2 / \beta_H$.

Let us now see what our polymer-type picture says about the nature of the degrees of freedom in the transition between the string and the black hole pictures. Consider the entropy S of the polymer string in $D = 4$ (the arguments are similar in other dimensions in which the above transition makes sense). At zero coupling, $S_0 \sim n$, the number of steps. This can be seen in the usual manner for a random walk by associating to each step a small whole number of independent possible directions (related to the number of space dimensions) p . The number of possible configurations gives the degeneracy $d \sim p^n$, so that the entropy $S = \ln d \sim n$.

In the intermediate range $g_0 < g < g_c$, the adiabatic increase of the coupling preserves the essential degrees of freedom associated with the string bits. Of course one no longer has a random walk, but up to a factor of order unity, the string bits retain most of their

degrees of freedom.

For a black hole whose mass is equal to the excited string state up to a factor of $O(1)$, $S_{BH} = A/4G \sim R_S^2/l_P^2$, where $l_P = gl_s$ is the Planck scale and $R_S \sim GM \sim l_P^2 M \sim l_P^2 n/l_s \sim g^2 nl_s$ is the Schwarzschild radius. We may then rewrite the BH entropy as

$$S_{BH} \sim \frac{n^2 l_P^2}{l_s^2} \sim n^2 g^2. \quad (11)$$

At the critical coupling $g_c \sim n^{-1/2}$, the entropy makes a smooth transition to the Bekenstein-Hawking area law form: $S_{BH} \sim n \sim S_0$. This makes sense, since we don't expect any violent changes in the entropy in the process of adiabatically increasing the coupling. It is interesting, however, that this entropy still represents a degeneracy of a polymer system with n steps. This can be seen as follows: at $g = g_c$, the size of the collapsed polymer string is given by $R \sim R_S \sim l_s$, or the size of one string bit, or one step. In order for n steps of size l_s to fit into a sphere of radius R_S , the number of possible positions to which each step can go must remain a small whole number, p' . Hence the degeneracy again has the random walk form $d \sim p'^n$. Of course, the polymer is no longer a random walk, but the degrees of freedom still remain essentially intact.

What happens in the black hole picture once the transition is complete? In this case, the area of the black hole is given by $A \sim R_S^2 \sim l_s^2 \sim (1/g^2)l_P^2 \sim nl_P^2$. So the horizon can be divided into n "pixels" each of area l_P^2 . Once the horizon forms, the degrees of freedom associated with it represent independent quantum states². Again, only a small whole number of possible states, q , is associated with each pixel, so that the total degeneracy is given by $d_{BH} \sim q^n$, with the entropy given by $S = \ln d \sim n \sim S_{BH}$. So essentially the random walk degrees of freedom turn into horizon surface degrees of freedom at the critical transition point. Another way of saying this is that the string bits project their information onto the horizon.

² The independence of the states is a result of the causally disconnected nature of the points on the horizon.

What happens if we continue increasing the coupling beyond the critical coupling g_c and into the black hole domain? Then the surface degrees of freedom should continue to account for the area law entropy. This can be seen in two ways: in the first way, we fix the mass while increasing g . This would necessarily no longer be an adiabatic change, and the entropy would have to change. In this case, $R_S \sim g^2 n l_s > l_s$ for $g > g_c = n^{-1/2}$, and the area $A \sim R^2 \sim g^4 n^2 l_s^2 \sim g^2 n^2 l_P^2$. The number of pixels is now given by $k = A/l_P^2 = g^2 n^2$, which is equal to Bekenstein-Hawking entropy, S_{BH} from (11).

The other way of extending beyond g_c is to continue adiabatically. In this case the entropy remains equal to the original random walk number n , but the mass changes so that $S \sim A/G \sim R_S^2/G \sim GM^2 \sim n$. This implies that $M^2 \sim n/l_P^2 \sim n/(g^2 l_s^2)$, so that $M \sim \sqrt{n}/(g l_s)$. It follows that $R_S \sim GM \sim S/M \sim n g l_s / \sqrt{n} = \sqrt{n} g l_s = \sqrt{n} l_P$, and the horizon area is $A \sim R_S^2 \sim n l_P^2$, representing n pixels and a quantum entropy of n .

The above results are in full agreement with expectations of the holographic principle, namely that the degrees of freedom of the black hole can be recovered through projection of the states onto the horizon surface. In conclusion, the above results not only independently confirm those of [4,8], but also suggest that the underlying degrees of freedom of quantum black holes in string theory remain associated with the original, stringy degrees of freedom. This further strengthens the string/black hole correspondence conjecture [1] by implying that in the transition to the strong-coupling limit, it is possible that the quantum string states somehow retain far more of their nature from the perturbative picture than might have been supposed. This idea, of course, requires further verification and more precise calculations, but nevertheless points to the robustness of the string/black hole correspondence principle.

References

- [1] L. Susskind, hep-th/9309145.
- [2] G. T. Horowitz and J. Polchinski, Phys. Rev. **D55** (1997) 6189.
- [3] J. Bekenstein, Lett. Nuov. Cimento **4** (1972) 737; Phys. Rev. **D7** (1973) 2333; Phys. Rev. **D9** (1974) 3292; S. W. Hawking, Nature **248** (1974) 30; Comm. Math. Phys. **43** (1975) 199.
- [4] G. T. Horowitz and J. Polchinski, Phys. Rev. **D57** (1998) 2557. See also S. Kalyana Rama, Phys. Lett. **B424** (1998) 39.
- [5] P. Salomonson and B. S. Skagerstam, Nucl. Phys. **B268** (1986) 349; Physica **A158** (1989) 499; D. Mitchell and N. Turok, Phys. Rev. Lett. **58** (1987) 1577; Nucl. Phys. **B294** (1987) 1138.
- [6] R. R. Khuri, Mod. Phys. Lett. **A13** (1998) 1407.
- [7] See C. B. Thorn, hep-th/9607204 and references therein; see also O. Bergman and C. B. Thorn, Nucl. Phys. **B502** (1997) 309.
- [8] T. Damour and G. Veneziano, hep-th/9907030.
- [9] See M. Doi and S. F. Edwards, *The Theory of Polymer Dynamics*, Clarendon Press, Oxford (1986) and references therein.
- [10] S. F. Edwards and M. Muthukumar, J. Chem. Phys. **89** (1988) 2435; S. F. Edwards and Y. Chen, J. Phys. **A21** (1988) 2963.